

The Effect Of Climate Change On The Communities Of Ogbaru Wetland Of South West Anambra State, Nigeria

Ezenwaji E.E.¹ Orji M.U.² Enete C.I.³ Otti, V. I.⁴

^{1,3}Department of Geography and Meteorology Nnamdi Azikiwe University, Awka, Nigeria.

²Department of Applied Microbiology and Brewing, Nnamdi Azikiwe University, Awka, Nigeria.

⁴Department of Civil Engineering Federal Polytechnic, Oko, Nigeria.

Abstract: - The aim of this study was to examine the effects of climate change on the livelihood of the inhabitants of the communities in Ogbaru wetland of Anambra State, Nigeria. A total of 300 questionnaires dealing on the effects of climate change in the 16 communities of the area were designed and served on the inhabitants through stratified sampling technique for 3 months May – July 2011. Also 40 years meteorological data on temperature, rainfall and relative humidity of the area were sourced from NIMET synoptic station in the area. Data were analysed with the use of Multiple Regression Technique to ascertain the collective contributions of the meteorological elements to climate change in the area. Furthermore, cluster analysis was employed to agglomerate the 16 communities into defined groups. Result shows that the three meteorological elements namely Temperature, Rainfall and Relative Humidity, altogether contributed 43% to the overall variation in climate change of the wetland while cluster analysis grouped the 16 communities into three, according to the degree of the impact of climate change on them. Measures aimed at adapting to the adverse impact of the change such as creation of earth dams around the farms, use of markings on walls to gauge the flood height of previous years, employing adequate construction techniques in road construction in the area etc. were discussed.

Keywords: - Climate, NIMET, agglomerate, meteorological, variation, synoptic.

I. INTRODUCTION

The effect of climate change to the global environment has dominated many international conferences, workshops and fora. Meanwhile studies of the effects of climate change have been carried out in many specific sectors of Nigeria. Aondo (1997), Ezenwaji (2010), and Nnaji (2011) investigated the influence of the phenomenon on water supply, while Adejuwon (2006) and SEI (2008) have all studied the effect of climate change on agricultural and food production. Various implications of climate change on socio-economic development of Nigeria have been assessed (Abatam, 2007; Ugwuanyi and Anekeje, 2009). Its effects on urban and rural environments have also been studied (Uyigwe and Agbo, 2007).

The natural resources of the Nigerian wetlands are indeed a major source of livelihood activity to wetlands inhabitants, while the nutrient rich soils are excellent for agricultural purposes. Climate variability and other human activities now threaten this unique habitat. Global temperature increases and causes the rise of the sea level, destroying coastal wetlands and drying up inland wetland basins that previously sustain human socio-economic activities. The environment of inland wetland which is the focus of this paper, has maintained biodiversity by providing critically important habitat to a wide range of wild life species. Flood protection is one important role which wetlands are known to play. Inland wetland soils act as natural sponges that trap and slowly release surface water. Wetland vegetation also helps to slow the speed of floodwaters and distribute them more gradually over the floodplain. The adverse impact of climate change on the agricultural and human activities of the wetland areas of the country need to be ameliorated through adaptive measures. Apart from the works of Ozor and Umehai (2010) and that of Ezenwaji (2010), the study of the effect of climate change in the Nigerian wetland is scanty. This paper, therefore, seeks to study the effect of climate change on the Ogbaru wetland of Anambra State and suggest adaptive management measures in dealing with the identified effects.

II. MATERIALS AND METHODS

Study Area

The entire Ogbaru Local Government Area which constitutes a large wetland zone is located in the south western part of Anambra State, and lies between latitudes 5°42' and 6°08'N and Longitudes 6°42' and 6°50'E. It is bound in the North by Onitsha South Local Government Area, in the east by Idemili South, Ekwusigo and Ihiala Local Government Areas, in the west by Delta State and in the south by Rivers and Imo States. The relief is a plain land of heights ranging from 0 – 50m and characterized by swampy conditions as a result of its alluvial mud content. Its geology is mainly alluvium while the river Niger, and Ulasiriver which is its

major tributary constitute the two major rivers in the area. However, there are local creeks and ponds all over its landscape. The vegetation is a mixture of fringing forests along the banks of the river Niger and guinea savannah in the hinterland. The climate is hot wet equatorial with average maximum temperature of 30°C and the average minimum of 24°C depending on the season of the year.




Figure 1: Map of Ogbaru Local Government Area.

Rainfall is experienced for 9 months of the year March – November, while dry season lasts from December to February (Monanu, 1975) with a total annual figure about 1,900mm. A range of agricultural and human activities take place here and they include fishing, lumbering, crop production, water and land transport etc. The area has a total population of according to the recent population figure from National Population Commission (2006). Sixteen riparian communities are located in the study area.

Data Collection

To ascertain the agricultural and human activities of the area affected by climate change, 300 questionnaires were designed and served on the inhabitants of the 16 communities that make up the area for 3 months (May – July) 2010, using stratified sampling technique with each community forming strata. Within each stratum, questionnaires were randomly distributed in proportion to the population of each community. In addition to the data collected from the questionnaires, the study also relied on temperature, rainfall and relative humidity data of the area for a period of 40 years, (1971 – 2010) sourced from the Nigerian Meteorological Agency, synoptic station, at Nkwelle, Onitsha, which is about 5kms to the study area. We took the data from this station as reliable because it is employed by government and other agencies to predict the meteorological elements for agricultural activities in the area.

Data Analysis

Data were analyzed with the use of Multiple Linear Regression to establish the contributions of the climate elements to the destruction of agricultural and human activities. The number of flood events in the area was used as the dependent variable, while the three climatic elements namely Temperature, Rainfall and relative Humidity were used as independent variable. (Table 1) Cluster Analytical Technique and Multiple Regression Analysis were employed in the study. The variables isolated from the questionnaire and from previous research in the area like Obidike(2006) were used to run cluster analysis and the result revealed a spatial arrangement of the effect of the variables on various communities in the area. Of all the methods of cluster analysis, discussed by Koo et-al (2005) was used in this study because it minimizes the loss of data information.

Results and Discussion

The 40 year average of temperature, rainfall and relative humidity of the area are presented in Table 1 while the labelling and explanation of the variables are seen in Table 2.

Table 1: 40 year 1971 – 2010 Average Temperature, Rainfall and Relative Humidity together with the number of major flood events in Ogbaru wetland

Year	Temp (°C)	Rainfall (mm)	Relative Humidity (%)	Numbers of major flood events *
1971	29.4	1662	44.4	6
1972	30.1	1612	49.2	4
1973	28.2	1566	39.2	2
1974	28.9	1633	46.9	4
1975	27.3	1687	58.4	6
1976	28.4	1849	56.8	5
1977	28.9	1877	58.1	8
1978	28.4	1820	62.3	8
1979	27.9	1829	54.1	6
1980	27.5	1864	59.3	6
1981	27.2	1842	52.1	4
1982	28.6	1872	60.1	5
1983	26.9	1828	58.9	5
1984	28.1	1794	54.6	3
1985	26.3	1849	58.7	4
1986	28.4	1812	65.4	9
1987	28.5	1756	63.4	7
1988	26.3	1791	64.1	6
1989	26.9	1920	65.9	8
1990	27.1	1846	64.1	7
1991	28.7	1821	64.8	4
1992	28.6	1920	65.7	4
1993	28.4	1872	62.6	3
1994	26.4	1856	68.3	4
1995	28.6	1824	67.2	5
1996	28.6	1881	66.4	2
1997	27.4	1866	67.9	3
1998	27.9	1846	67.8	4
1999	27.4	1822	67.4	4
2000	27.9	1814	68.3	4
2001	28.7	1869	67.3	2
2002	28.6	1881	67.4	3

2003	28.1	1826	64.2	3
2004	28.4	1883	65.1	5
2005	27.3	1842	62.4	4
2006	27.9	1809	64.1	3
2007	27.7	1832	61.2	5
2008	27.4	1881	66.2	6
2009	28.2	1845	67.2	6
2010	28.3	1861	68.1	7

Source: Nigerian Meteorological Agency Amawbia Synoptic Station, (2011).

* Obidike (2006) and field work (2010).

Table 2: Parametisation of Variables of Climate Change Effects in OgaruWetland

Variable label	Explanation
SWELL	Number of wells that swell up because of flooding in 2010
AGRIC	Number of Agric farms that were destroyed by flood in 2010
LOGG	Number of locations that were waterlogged for the entire rainy season
POLL	Number of water sources that were polluted with microbial contaminants from floods in 2010
SILT	Number of streams that were silted in 2010
DISP	Number of families displaced in 2010 as a result of flooding
FISH	Average price of fishes harvested in August 2010
DES	Number of roads destroyed by flooding and soil erosion in 2010

Also Table 3 shows the data gathered from the above variables during the field work.

Table 3: Field Data of the Resultant Effect of the Three Climatic Elements

S/N	Community	SWELL	AGRIC	LOGG	POLL	SILT	DISP	FISH	DES
1	Okpoko	64	0	10	15	10	0	0	11
2	Odekpe	32	4	16	14	8	6	600	2
3	Ohuta	10	3	8	8	6	9	1000	2
4	Atani	15	10	18	12	10	10	1000	2
5	AkiliOzizor	20	6	9	14	14	12	1500	3
6	Umu-Uzu	6	4	3	10	11	8	1500	2
7	Ochuche	4	8	10	5	10	8	1500	3
8	Ogbakuba	9	6	18	4	8	7	800	2
9	Umunankwo	4	11	12	7	6	10	600	2
10	Osomala	6	15	22	5	13	12	800	2
11	Obagwe	4	13	6	2	4	6	1000	2
12	Oguaniocha	8	22	16	10	10	8	1000	4
13	AkiliOgidi	7	29	20	5	21	29	600	2
14	OguIkpele	4	18	20	4	10	28	2500	3
15	Mputu	6	4	8	3	6	4	600	2
16	Umuodu	3	6	4	5	8	4	600	3

Data in Table 1 were analyzed using the Multiple Regression Analysis to determine the contributions of the climatic elements to the destruction of the agricultural and human activities of the area. The climatic elements were labelled as follows;

Variable	Label
Temperature	TEMP
Rainfall	RAIN
Relative Humidity	REHU

When the 3 climate elements (used as independent variables) were correlated on the number of flood events, (the dependent variable)it produced the result shown in Table 5.

Table 5: Result of the Correlation Analysis

Variables	Flood	Temp.	Rainfall	Humidity
Flood	1.000	- 0.74	- 0.108	0.085
Temp.	- 0.074	1.000	- 0.318	- 0.275
Rainfall	0.108	- 0.318	1.000	0.757*
Humidity	0.085	- 0.275	0.757*	1.000

* High correlation coefficients.

The correlation in Table 5 is generally low except for rainfall and atmospheric humidity. This is the natural trend as the higher the rainfall the more likely is relative humidity. Conversely, flood and all the parameters are weakly correlated but their various relationships are expected. For example the negative relationship between flood and temperature means that as temperature is reducing, flood is increasing because within that situation there are chances of increased rainfall. The result of the Regression Analysis (Table 6) indicates that the combined atmospheric parameters contributed 43% to flooding in the area. This means that anthropogenic and environmental factors take up 57% of the flooding in the wetland. This is expected because of the influence of large body of water which supplies ample moisture during evaporation which eventually comes back to the environment as rain which floods the environment.

Table 6: Result of the Regression Analysis

Model	R	R ²	Std. Error of estimate
1	116 ^a	0.043	1.837

A predictors: (constant) Humidity, Temp, Rainfall.

The ANOVA performed to ascertain the strength of our regression shows that its p-level (sig) is 0.03 which is below 0.05 level (Table 7). This indicates that the regression model is strong in interpreting the relationship between Flood and Climatic parameters.

Table 7: ANOVA Table

Model	Sum of squares	Df	Mean square	F	Sig
Regression	1.646	3	0.549	0.163	0.03
Residual	121.454	36	3.374		
Total	123.100	39			

However, based on the field data generated for each of the 8 variables a cluster analytical method earlier employed by Koo *et-al*(2005) was performed and the result shows the spatial impact of climate change on the 16 communities as produced by Cluster Analysis model in Figure 2, Table 8 however, shows the regionalization of the impact of climate change in the communities located within Ogbaru wetland.

Dendrogram




Fig. 2: The Dendrogram of the area.

From this figure, the area was organized by the model in a three group cluster as follows;

Table 8: Climatic Change impact similarities of Communities in Ogbaru

Groups	Group features	Community and their identification number
1.	High climatic change impact	Okpoko (1) and Oguikpele (14)
2.	Moderate climate change impact	Odekpe (2) and AkiliOgidi (13)
3.	Low climatic change impact	Ohital (3), Atani (4), AkiliOzizor (5), Umuzu (6), Ochuche (7), Ogbakuba (8), Umunankwo (9), Ossamala (10), Obegwe (11), Ogwuanocha (12), Mputu (15) and Umuodu (16).

The cluster analysis has thus agglomerated the 16 communities into 3 groups based on the degrees of impact of climate change on them, viz;

Group 1 communities are those that are flooded for over 8 months of the year. Included here are Okpoko which in the urban area of the wetland. The high level of impervious covering, blocking of drainage systems and manifest absence of any type of vegetation in this area together with its very low lying relief may have accounted for this. Then Oguikpele which is another community in this group is also usually overran by flood for mostpart of the year at times up to 9 months. The main reason here is as a result of its low lying relief and its location at the point where the river Niger seem to have one of the greatest discharge rates. The overflow of the river bank by the water from the river into the overlying plain is therefore an essential geographical feature of this location.

Group 2 are two communities that have moderate climate change impact. As in group 1, here only two communities are included and they are Odekpe which has a considerable level of urbanization. The other community is AkiliOgidi which is very close to Oguikpele. All the factors that gave rise to high level of flood in okpoko and Oguikpele also account for the situation in these two communities.

Group 3 are 12 communities that are classified as areas with low climate change impact. These are Ohita, Atani, AkiliOzizor, Umuzu, Ochuche, Ogbakuba, Umunankwo, Ossomala, Obegwe, Ogwuanocha, Mputu and Umuodu. Half of the numbers of communities in this group are located at the river bank while the remaining half are located away from the banks. Those that are not located at the banks are Ohita, Atani, Umuzu, Umuodu, Mputu and Ogwuanochawhich always have less severe flood problems.

It could be indicated here that a whole range of agricultural and human activities are seriously affected by the flooding activities in the area. The generally low-lying nature of Ogbaru wetland makes it prone to flooding which destroy farmland, sack communities, cut road network, pollute rivers and water sources, reduce fish harvest and a number of other havocs. The following adaptation measures are suggested.

1. The inhabitants of the area have been practicing adaptation measures for ages which should now be urgently strengthened. From our fieldwork investigation, we observed that so many farms had earth dykes of small heights around their farms. Our enquiry revealed that they are built to avoid inflow of flood waters into their farms. In this regard, government should encourage the farmers with seed funds to help them continue with this practice.
2. The inhabitants make markings on the walls at the banks of river Niger to gauge the height of flood of the previous year and to know when a succeeding years flood has gone beyond that height, the inhabitation usual create marks at the banks of the river. This is indeed a warning signal to alert them as to when to move out of an area to another safe location. Although, this method seems crude, it, however, calls for an urgent action from relevant government agencies to increase efforts on flood forecasting in the area so as to alert the local people whenever there is an impending flood.
3. Again, because fishing activities are hampered by high volume of floods, the local people have been creating artificial fish holes in their farms from where they harvest fishes when floods recede usually at the end of flood episodes. During floods, fishes move into such holes as ready homes and fish farmers go there to harvest large quantities when the floods recede. In this regard, the government can help by creating artificial pondages at the flood plain where the fishes could find sanctuary during inclement flood period and harvest thereafter.
4. Recent road construction methods have not addressed the peculiar geographic nature of the area, leading to the incessant collapse of both tared and earth roads. It is suggested that any road construction of the area should ensure that enough stone base and drainages are provided to ensure that flood waters do not destroy them.

5. Also in the part of the wetland that is already urbanized, all human activities that aid flooding should be avoided. In Okpoko area of the wetland, the level of flooding whenever it rains is frightening which calls for immediate urban planning measures.

III. CONCLUSION

Since a range of damages from future flood of the area is inevitable, efforts should be made to promote the adaptation measures outlined in this paper. It is indeed evident that the old adaptation strategies are no longer coping with the increasing menace of climate change in the area. Modern measures should therefore be adopted. The efforts of the State government and indeed other government agencies should be concentrated on areas with high flood prone risks such as Okpoko and Oguikpele. Finally it is envisaged that since we have achieved a categorization of degree of climate change impact in the area, all adaptation measures should be in response to the extent of the impact on the communities.

REFERENCES

- [1] Abatam, A.A. (2007): *Effect of Air Pollution on the Socio-Economic Development in Nigeria* <http://ihy2007.org/img/IHY>, Nigeria Abatam.doc
- [2] Adejuwon, I.O. (2006): "Food Crop Production in Nigeria II: Potential effects of climate change". *Climatic Research*, Vol. 32, pp 229-245.
- [3] Anpover, A.T. (1997): *Droughts, Rainfall and Rural Water Supply in Northern Nigeria*. PhD Thesis, Department of Geography, McMaster University.
- [4] Ezenwaji, A.N. (2010). *Flood Mitigation Parameters of Apiti Wetland Soil of Umuawulu, Awka South L.G.A., Anambra State*. An Unpublished M.Sc. Thesis, Department of Zoology, NnamdiAzikiwe University, Awka.
- [5] Ezenwaji, E.E. (2010): "Climate Change and Household Water Supply Shortages in Onitsha Urban Area, Nigeria" in AnyadikeR.N.C., Madu, I.A. and Ajaero, C.K. (eds), *Climate Change and the Nigerian Environment*, Jamoe Publishers, Enugu.
- [6] Koo, J.Y., Yu, M.J., Kim, S.G., Shim, M.H., & Koizumi, A. (2005): "Estimating Regional Water Demand in Seoul, South Korea, Using Principal Compound Analysis and Cluster Analysis". *Water Science and Technology: Water Supply*, Vol. 5, No. 1, pp 1-7.
- [7] Monanu, P.C. (1975): "Rainfall" in OfomataG.E.K. (ed) *Nigeria in Maps: Eastern States*. Ethiopia Press Benin City.
- [8] National Population Commission (2006): *2006 Local Government Area Population*. Govt. Printer Abuja.
- [9] Nnaji, A.O. (2011): "Implications of Recent Rainfall Variability for urban Water Supply and Sustenance in Owerri, Imo State". Paper delivered at the 1st National Workshop of the Environment Development Initiative for Africa (edia) Awka, 24th – 26th August, 2011.
- [10] Obidike, J. (2006). The Development of Ogbaru Area of Anambra State. *Journal of Regional Development*, Vol. 4, No. 3, pp 60 -71.
- [11] Ozor, N. and Umehai, M.C. (2010): Effects of Climate Change on the Livelihoods of wetlands inhabitants in Nigeria. A Review, in AnyadikeR.N.C., Madu, I.A., and Ajaero, C.K. (eds.) *Climate Change and the Nigerian Environment*.
- [12] SEI (2008): *Climate Change and Adaptation to African Agriculture*. Report prepared by Rockefeller Foundation by Stockholm Environment Initiative, March 2008, pp 54.
- [13] Ugwuany, A.C. and Anekeje, B. (2009): "Increasing Climate Change effects on Humans Activities of some rural communities in Benue area, Nigeria". *Journal of Continental Ecology*, Vol. 2, No. 2, pp 75-87.
- [14] Uyigue, E. and Agbo, M. (2007): "Coping with Climate Change and Environmental Degradation in the Niger Delta of Southern Nigeria. *Community Research and Development Centre, Nigeria CREDC*.