

## On Some Double Integral Transform of Aleph $\aleph$ -Function

Yashwant Singh

*Department of Mathematics  
 Government College, Kaladera, Jaipur (Rajasthan), India  
 Corresponding Author: Yashwant Singh*

**Abstract:** In the present paper , the author will establish a double integral transform of Aleph  $\aleph$ -function which leads to yet another interesting process of augmenting the parameters in the  $\aleph$ -function. The result is of general character and on specializing the parameters suitably yields several interesting results as particular cases.

**Keywords:**  $\aleph$ -function, Euler Transformation,  $H$ -function, Hypergeometric Function, Integral Transformation.

(2010 Mathematics subject classification: 33C99)

Date of Submission: 04-01-2019

Date of acceptance: 19-01-2019

### I. INTRODUCTION

Rainville [6, p.104], Abdul Halim and Al-Salam [1] have shown that the single and double Euler transformations of the hypergeometric function  ${}_pF_q$  are effective tools for augmenting its parameters. Srivastava and Singhal [9] and Srivastava and Joshi [10] have discussed some similar interesting properties of  ${}_pF_q$  in double  $H$ -function and double Whittaker transforms respectively.

In what follows for the sake of brevity, we have used the symbols  $(a_r, \alpha_r), \Delta(r, a), \Delta(r, \pm a), \Delta((r, a_p))$  to denote the set of parameters  $(a_1, \alpha_1), \dots, (a_r, \alpha_r); \frac{a}{r}, \frac{a+1}{r}, \dots, \frac{a+r-1}{r}; \Delta(r, a), \Delta(r, -a)$  and  $\Delta(r, a_1), \Delta(r, a_2), \dots, \Delta(r, a_p)$  respectively.

The  $\aleph$ -function introduced by Suland et.al. [11] defined and represented in the following form:

$$\begin{aligned} \aleph[z] &= \aleph_{p_i, q_i; \tau_i; r}^{m, n}[z] = \aleph_{p_i, q_i; \tau_i; r}^{m, n} \left[ z \mid \begin{array}{l} (a_j, \alpha_j)_{1, n}, [\tau_i(a_{ji}, \alpha_{ji})]_{n+1, p_i} \\ (b_j, \beta_j)_{1, m}, [\tau_i(b_{ji}, \beta_{ji})]_{m+1, q_i} \end{array} \right] \\ &= \frac{1}{2\pi\omega} \int_L \theta(s) z^s ds \end{aligned} \quad (1.1)$$

Where  $\omega = \sqrt{-1}$ ;

$$\theta(s) = \frac{\prod_{j=1}^m \Gamma(b_j - \beta_j s) \prod_{j=1}^n \Gamma(1 - a_j + \alpha_j s)}{\sum_{i=1}^r \tau_i \left\{ \prod_{j=m+1}^{q_i} \Gamma(1 - b_{ji} + \beta_{ji} s) \prod_{j=n+1}^{p_i} \Gamma(a_{ji} - \alpha_{ji} s) \right\}} \quad (1.2)$$

We shall use the following notation:

$$\begin{aligned} A^* &= (a_j, \alpha_j)_{1, n}, [\tau_i(a_{ji}, \alpha_{ji})]_{n+1, p_i}, B^* = (b_j, \beta_j)_{1, m}, [\tau_i(b_{ji}, \beta_{ji})]_{m+1, q_i} \\ C^* &= (a_j, \alpha_j)_{1, f}, [\tau_i(a_{ji}, \alpha_{ji})]_{f+1, u_i}, D^* = (b_j, \beta_j)_{1, g}, [\tau_i(b_{ji}, \beta_{ji})]_{g+1, v_i} \end{aligned}$$

## II. MAIN RESULT

In this section, we have established the following double integral transform of  $\aleph$ -function:

If  $s, k$  and  $r$  are positive integers, then

$$\begin{aligned} \int_0^\infty \int_0^\infty x^{\alpha-1} y^{\beta-1} (x+y)^\sigma \aleph_{u_i, v_i; \tau_i; r}^{f, g} \left[ \lambda(x+y) \Big|_{D^*}^{C^*} \right] \aleph_{p_i, q_i; \tau_i; r}^{m, n} \left[ t x^s y^k (x+y)^r \Big|_{B^*}^{A^*} \right] dx dy = \\ (2\pi)^{(1-D)\left(f+g-\frac{1}{2}u-\frac{1}{2}v\right)+\frac{1}{2}} D^{\sum_i^v d_j - \sum_i^u c_j + \left(A-\frac{1}{2}\right)(u-v)} \frac{s^{\alpha-\frac{1}{2}} k^{\beta-\frac{1}{2}}}{\lambda^{\alpha+\beta+\sigma} (s+k)^{\alpha+\beta-\frac{1}{2}}} \\ \aleph_{p_i+\rho+Dv_i, q_i+\rho+Du_i; \tau_i; r}^{m+Dg, n+\rho+Df} \left[ \frac{t \delta D^{D(v-u)}}{\lambda^D} \Big|_{\Delta((D, 1-A-D_f)), \Delta(s, 1-\alpha), \Delta(k, 1-\beta), A^*, \Delta(D, 1-A-d_{f+1}), \dots, \Delta(D, 1-A-d_u)}^{\Delta((D, 1-A-C_g)), B^*, \Delta(k+s, 1-\alpha-\beta), \Delta(D, 1-A-c_{g+1}), \dots, \Delta(D, 1-A-c_v)} \right], \quad (2.1) \end{aligned}$$

Where  $\gamma = \frac{s^s k^k}{(s+k)^{s+k}}$ ,  $\rho = s+k$ ,  $D = s+k+r$ ,  $A = \alpha + \beta + \sigma$ ,

$0 \leq Dg \leq Du \leq Dv < Du + q - p, u + v - 2g \leq 2f \leq 2v, 0 \leq n \leq p, p + q - 2n < 2m \leq 2q$ ,

$$\operatorname{Re} \left( \min \tau_i \frac{d_{ji}}{\delta_{ji}} + D \min \tau_i \frac{b_{ji}}{\beta_{ji}} \right) > \operatorname{Re}(-A) > \operatorname{Re} \left[ D \left( \frac{s-\alpha}{s}, \frac{k-\beta}{k}, a_l \right) + C_t - D - 1 \right]$$

$i = 1, 2, \dots, f; j = 1, 2, \dots, m; l = 1, 2, \dots, n; t = 1, 2, \dots, g; u, \operatorname{Re}(\min C_i + A) - v$ ,

$$\operatorname{Re} \left( \max \frac{d_j}{\delta_j} + A \right) - uD + v + \frac{1}{2}D(Dv - Du + 1) > D(Dv - Du), \operatorname{Re} \max \left( \frac{s-\alpha}{s}, \frac{k-\beta}{k}, a_l \right),$$

$i = 1, 2, \dots, u; j = 1, 2, \dots, v; l = 1, 2, \dots, u; |\arg \lambda| \leq \left( f + g - \frac{1}{2}u - \frac{1}{2}v \right) \pi$ ,

$|\arg t| < \left( m + n - \frac{1}{2}p_i - \frac{1}{2}q_i \right) \pi, \operatorname{Re} \left( \alpha + s\tau_i \frac{b_{ji}}{\beta_{ji}} \right) > 0, \operatorname{Re} \left( \beta + k\tau_i \frac{b_{ji}}{\beta_{ji}} \right) > 0, j = 1, 2, \dots, m$

And the double integral converges.

Proof: To prove (2.1), we start with the following known result [2, p. 177]

$$\int_0^\infty \int_0^\infty \phi(x+y) x^{\alpha-1} y^{\beta-1} dx dy = B(\alpha, \beta) \int_0^\infty \phi(z) z^{\alpha+\beta-1} dz \quad (2.2)$$

Which is valid for  $\operatorname{Re}(\alpha) > 0$  and  $\operatorname{Re}(\beta) > 0$ .

It is easy to prove by following the technique of reversing the order of integrations, that

$$\begin{aligned} \int_0^\infty \int_0^\infty \phi(x+y) x^{\alpha-1} y^{\beta-1} \aleph_{p_i, q_i; \tau_i; r}^{m, n} \left[ t x^s y^k (x+y)^r \Big|_{B^*}^{A^*} \right] dx dy = \sqrt{2\pi} \frac{s^{\alpha-\frac{1}{2}} k^{\beta-\frac{1}{2}}}{(s+k)^{\alpha+\beta-\frac{1}{2}}} \\ \int_0^\infty \phi(z) z^{\alpha+\beta-1} \aleph_{p_i+\rho, q_i+\rho; \tau_i; r}^{m+n+\rho} \left[ t \delta z^D \Big|_{B^*, \Delta(k+s, 1-\alpha-\beta)}^{\Delta(s, 1-\alpha), \Delta(k, 1-\beta), A^*} \right] dz \quad (2.3) \end{aligned}$$

Where  $s, k$  and  $r$  are positive integers,

$$\begin{aligned} \delta = \frac{s^s k^k}{(s+k)^{s+k}}, \rho = s+k, D = s+k+r, p+q < 2(m+n), |\arg t| < \left( m + n - \frac{1}{2}p_i - \frac{1}{2}q_i \right) \pi, \\ \operatorname{Re} \left( \alpha + s\tau_i \frac{b_{ji}}{\beta_{ji}} \right) > 0, \operatorname{Re} \left( \beta + k\tau_i \frac{b_{ji}}{\beta_{ji}} \right) > 0, j = 1, 2, \dots, m. \end{aligned}$$

In (2.3), taking

$$\phi(z) = z^\sigma \aleph_{u_i, v_i; \tau_i; r}^{f, g} \left[ \lambda z \Big|_{D^*}^{C^*} \right]$$

And evaluating the integral on the right hand side using [8, p.401] the result (2.1) follows.

### III. PARTICULAR CASES

On choosing the parameters suitably in (2.1), several known and unknown results are obtained as particular cases. However, we mention some of the interesting results here.

(a)

Taking

$$f = v_i = 2, g = 0, u = 1, c_1 = \frac{1}{2}, d_1 = v, d_2 = -v, \sigma = \mu + \frac{1}{2}, \alpha_j = \beta_j = \delta_j = \gamma_j = 1, \tau_i = 1, r = 1 \text{ in (2.1)}$$

and using [3, p.216, (5)]

$$H_{1,2}^{2,0} \left[ x \Big|_{(b,1),(-b,1)}^{\left(\frac{1}{2},1\right)} \right] = \pi^{-\frac{1}{2}} e^{-\frac{1}{2}x} K_b \left( -\frac{1}{2}x \right),$$

We obtain

$$\begin{aligned} & \int_0^\infty \int_0^\infty x^{\alpha-1} y^{\beta-1} (x+y)^{\mu+\frac{1}{2}} a^{-\frac{1}{2}\lambda(x+y)} K_\nu \left\{ -\frac{1}{2} \lambda(x+y) \right\} H_{p,q}^{m,n} \left[ tx^s y^k (x+y)^r \Big|_{B^*}^{A^*} \right] dx dy = \\ & (2\pi)^{-\frac{1}{2}(2-D)\left(f+g-\frac{1}{2}u-\frac{1}{2}v\right)+\frac{1}{2}} D^{A-1} \sqrt{\pi} \frac{s^{\alpha-\frac{1}{2}} k^{\beta-\frac{1}{2}}}{\lambda^{\alpha+\beta+\sigma} (s+k)^{\alpha+\beta-\frac{1}{2}}} \\ & H_{p+\rho+2D, q+\rho+D}^{m,n+\rho+2D} \left[ \frac{t\delta D^D}{\lambda^D} \Big|_{\Delta((D,1-A\mp\nu)), \Delta(s,1-\alpha), \Delta(k,1-\beta), A^*}^{\Delta((D,1-A\mp\nu)), \Delta(s,1-\alpha), \Delta(k,1-\beta), A^*} \right], \end{aligned} \quad (3.1)$$

Where  $\delta, D$  and  $\lambda$  have the same value as (2.1) and

$$A = \mu + \alpha + \beta + \frac{1}{2}; p + q < 2(m+n), \operatorname{Re}(\alpha + s \frac{b_j}{\beta_j} \pm \nu) > 0, \operatorname{Re}(\beta + s \frac{b_j}{\beta_j} \pm \nu) > 0,$$

$$\operatorname{Re} \left( \alpha + \beta + \mu \pm \nu + D \frac{b_j}{\beta_j} + \frac{1}{2} \right) > 0, j = 1, 2, \dots, m; \operatorname{Re}(\lambda) > 0, |\arg t| < \left( m + n - \frac{1}{2}p - \frac{1}{2}q \right) \pi.$$

(b) Further, replacing  $q, t$  and  $(a_p, \alpha_p)$  by  $q+1, -t$  and  $(1-a_p, \alpha_p)$  respectively and then putting  $m=1, n=p, b_1=0, b_{j+1}=1-b_j$  ( $j=1, 2, \dots, q$ ), using the result [3,p. 215, (1)] and [3,p. 4, (11)], we obtain an interesting result obtained by Srivastava and Singhal [9]:

$$\begin{aligned} & \int_0^\infty \int_0^\infty x^{\alpha-1} y^{\beta-1} (x+y)^{\mu+\frac{1}{2}} a^{-\frac{1}{2}\lambda(x+y)} K_\nu \left\{ -\frac{1}{2} \lambda(x+y) \right\} {}_p F_q \left[ \begin{matrix} (a_p, \alpha_p) \\ (b_q, \beta_q) \end{matrix} ; tx^s y^k (x+y)^r \right] dx dy = \\ & \frac{\sqrt{\pi} \Gamma \left( \alpha + \beta + \mu \pm \nu + \frac{1}{2} \right)}{\lambda^{\alpha+\beta+\mu+\frac{1}{2}} \Gamma(\alpha + \beta + \mu + 1)} B(\alpha, \beta) \\ & {}_p F_q \left[ t \delta \left( \frac{s+k+r}{\lambda} \right)^{s+k+r} \Big|_{(b_q, 1), \Delta(s+k, \alpha+\beta), \Delta(k+s, \alpha+\beta+\mu+1)}^{\Delta(s+k+r, \alpha+\beta+\mu \pm \nu + \frac{1}{2}), \Delta(s, \alpha), \Delta(k, \beta), (a_p, 1)} \right], \end{aligned} \quad (3.2)$$

provided  $\operatorname{Re}(\mu + \alpha + \beta \pm \nu + \frac{1}{2}) > 0, \operatorname{Re}(\alpha) > 0, \operatorname{Re}(\beta) > 0$ .

(c) Setting  $v = f = 2, g = 0, u = 1, c_1 = 1 - \mu, d_1 = \frac{1}{2} + v, d_2 = \frac{1}{2} - v, \tau_i = 1, r = 1$  in (2.1) and using the known formula [3,p.216,(6)]

$$H_{1,2}^{2,0} \left[ x \left| \begin{smallmatrix} (1-k,1) \\ (\frac{1}{2}+m,1), (\frac{1}{2}-m,1) \end{smallmatrix} \right. \right] = e^{-\frac{1}{2}x} W_{k,m}(x),$$

We have

$$\begin{aligned} & \int_0^\infty \int_0^\infty x^{\alpha-1} y^{\beta-1} (x+y)^\sigma e^{-\frac{1}{2}\lambda(x+y)} W_{\mu,v}[\lambda(x+y)] H_{p,q}^{m,n} \left[ tx^s y^k (x+y)^r \left| \begin{smallmatrix} A^* \\ B^* \end{smallmatrix} \right. \right] dx dy = \\ & (2\pi)^{\frac{1}{2}(2-D)} D^{\mu+A-\frac{1}{2}} \frac{s^{\frac{\alpha-1}{2}} k^{\frac{\beta-1}{2}}}{\lambda^{\alpha+\beta+\sigma} (s+k)^{\frac{\alpha+\beta-1}{2}}} \\ & H_{p+\rho+2D, q+\rho+D}^{m,n+\rho+2D} \left[ \frac{t\delta D^D}{\lambda^D} \left| \begin{smallmatrix} \Delta(D, \frac{1}{2}-A\pm v), \Delta(s, 1-\alpha), \Delta(k, 1-\beta), A^* \\ B^*, \Delta(k+s, 1-\alpha-\beta), \Delta(D, \mu-A) \end{smallmatrix} \right. \right], \end{aligned} \quad (3.3)$$

Where

$D, \rho, \delta$  and  $A$  are given in (2.1);

$$\begin{aligned} p+q < 2(m+n), |\arg t| < \left( m+n - \frac{1}{2}p - \frac{1}{2}q \right)\pi, \operatorname{Re}(\lambda) > 0, \operatorname{Re}(\alpha + sb_j) > 0, \\ \operatorname{Re}(k + sb_j) > 0, \operatorname{Re} \left( m+n+\sigma+Db_j \pm v + \frac{1}{2} \right) > 0, j=1, 2, \dots, m. \end{aligned}$$

(d) Further, replacing  $q, t$  and  $(a_p, \alpha_p)$  by  $q+1, -t$  and  $(1-a_p, \alpha_p)$  respectively and then putting  $m=1, n=p, b_1=0, b_{j+1}=1-b_j (j=1, 2, \dots, q)$  and using the result [p.215,(1)], (3.3) reduces to a result due to Srivastava and Joshi [10,p.19,(2.3)]

$$\begin{aligned} & \int_0^\infty \int_0^\infty x^{\alpha-1} y^{\beta-1} (x+y)^\sigma e^{-\frac{1}{2}\lambda(x+y)} W_{\mu,v} \{ \lambda(x+y) \} {}_p F_q \left[ \begin{smallmatrix} (a_p, \alpha_p) \\ (b_q, \beta_q) \end{smallmatrix} : tx^s y^k (x+y)^r \right] dx dy = \\ & \frac{\Gamma \left( \alpha + \beta + \sigma \pm v + \frac{1}{2} \right)}{\lambda^{\alpha+\beta+\sigma} \Gamma(\alpha + \beta + \sigma - \mu + 1)} B(\alpha, \beta) \\ & {}_{p+3s+3k+2r} F_{q+2s+2k+r} \left[ t\delta \delta' \left| \begin{smallmatrix} \Delta(s+k+r, \alpha+\beta+\sigma \pm v + \frac{1}{2}), \Delta(s, \alpha), \Delta(k, \beta), (a_p, 1) \\ (b_q, 1), \Delta(s+k, \alpha+\beta), \Delta(k+s+r, \alpha+\beta+\sigma-\mu+1) \end{smallmatrix} \right. \right] \end{aligned} \quad (3.4)$$

Where

$$\delta = \frac{s^s k^k}{(s+k)^{s+k}}, \delta' = \left( \frac{s+k+r}{\lambda} \right)^{s+k+r}$$

$\operatorname{Re}(\alpha) > 0, \operatorname{Re}(\beta) > 0, \operatorname{Re}(\lambda) > 0, \operatorname{Re} \left( \alpha + \beta + \sigma \pm v + \frac{1}{2} \right) > 0$  and the resulting hypergeometric series converges.

With  $\mu=0, v=\pm\frac{1}{2}$  and  $\sigma=-\frac{1}{2}$ , (3.4) reduces to the earlier results of Jain [5] and Singh [7].

(e)

Choosing

$$f=g=u=1, v=2, c_1=1-k, d_1=\frac{1}{2}+M, d_2=\frac{1}{2}-M, \alpha_j=\beta_j=\delta_j=\gamma_j=1, \tau_i=1, r=1 \text{ in } (2.1)$$

and using the known result

$$H_{1,2}^{1,1} \left[ x \begin{matrix} (1-k,1) \\ (\frac{1}{2}+m,1), (\frac{1}{2}-m,1) \end{matrix} \right] = \frac{\Gamma\left(\frac{1}{2}+k+m\right)}{\Gamma(2m+1)} e^{-\frac{1}{2}x} M_{k,m}(x),$$

We obtain

$$\begin{aligned} & \int_0^\infty \int_0^\infty x^{\alpha-1} y^{\beta-1} (x+y)^\sigma e^{-\frac{1}{2}\lambda(x+y)} M_{k,m}[\lambda(x+y)] H_{p,q}^{m,n} \left[ tx^s y^k (x+y)^r \Big| {}_{B^*}^{A^*} \right] dx dy = \\ & (2\pi)^{\frac{1}{2}(2-D)} D^{k+A-\frac{1}{2}} \frac{\Gamma(2m+1)}{\Gamma\left(k+m+\frac{1}{2}\right)} \frac{s^{\alpha-\frac{1}{2}} k^{\beta-\frac{1}{2}}}{\lambda^{\alpha+\beta+\sigma} (s+k)^{\alpha+\beta-\frac{1}{2}}} \\ & H_{p+\rho+2D, q+\rho+D}^{m+D, n+\rho+D} \left[ \frac{t\delta D^D}{\lambda^D} \begin{matrix} \Delta(D, \frac{1}{2}-A-m), \Delta(s, 1-\alpha), \Delta(k, 1-\beta), \Delta(D, \frac{1}{2}-A+m) \\ \Delta(k+s, 1-\alpha-\beta), \Delta(D, k-A) \end{matrix} \right], \end{aligned} \quad (3.5)$$

Where

$D, \rho, \delta$  and  $A$  are given in (2.1);

$$\begin{aligned} p+q < 2(m+n), |\arg t| < \left( m+n - \frac{1}{2}p - \frac{1}{2}q \right) \pi, \operatorname{Re}(\lambda) > 0, \operatorname{Re}(\alpha + sb_j) > 0, \\ \operatorname{Re}(\beta + kb_j) > 0, \operatorname{Re}\left(\alpha + \beta + \sigma + Db_j + m + \frac{1}{2}\right) > 0, j = 1, 2, \dots, m. \end{aligned}$$

(f) Substituting  $f = 1, g = u = 0, v = 2, d_1 = \frac{1}{2}v, d_2 = -\frac{1}{2}v, \tau_i = 1, r = 1$  and using the result [3,p.216,(3)]

$$H_{0,2}^{1,0} \left[ x \begin{matrix} - \\ \left( \frac{1}{2}v, 1 \right), \left( -\frac{1}{2}v, 1 \right) \end{matrix} \right] = J_v(2\sqrt{x}),$$

(2.1) reduces to

$$\begin{aligned} & \int_0^\infty \int_0^\infty x^{\alpha-1} y^{\beta-1} (x+y)^\sigma J_v(2\sqrt{\lambda(x+y)}) H_{p,q}^{m,n} \left[ tx^s y^k \Big| {}_{B^*}^{A^*} \right] dx dy \\ & = \sqrt{2\pi} \frac{D^{2A-1} s^{\alpha-\frac{1}{2}} k^{\beta-\frac{1}{2}}}{\lambda^{\alpha+\beta+\sigma} (s+k)^{\alpha+\beta-\frac{1}{2}}} \\ & H_{p+\rho+2D, q+p}^{m,n+\rho+D} \left[ t\delta \left( \frac{D}{\lambda} \right)^D \begin{matrix} \Delta\left(D, 1-A-\frac{1}{2}v\right), \Delta(s, 1-\alpha), \Delta(k, 1-\beta), A^*, \Delta\left(D, 1-A+\frac{1}{2}v\right) \\ B^*, \Delta(k+s, 1-\alpha-\beta) \end{matrix} \right] \end{aligned} \quad (3.6)$$

Where  $\delta, D, \rho$  and  $A$  have the same values given in (2.1);

$$\begin{aligned} p+q < 2(m+n), |\arg t| < \left( m+n - \frac{1}{2}p - \frac{1}{2}q \right) \pi, \operatorname{Re}(\lambda) > 0, \operatorname{Re}(\alpha + sb_j) > 0, \\ \operatorname{Re}(\beta + kb_j) > 0, \operatorname{Re}\left(\alpha + \beta + \sigma + \frac{1}{2}v + Db_j\right) > 0, j = 1, 2, \dots, m; \\ \operatorname{Re}(\alpha + \beta + \sigma - D + Da_i), \frac{1}{4}, i = 1, 2, \dots, n. \end{aligned}$$

In view of the numerous properties of Aleph ( $\aleph$ ) -function, on specializing the parameters suitably, a large number of interesting results may be obtained as particular case.

**REFERENCES**

- [1]. Abdul-Halim,N. and Al-Salam, W.A.; Double Euler transformation of certain hypergeometric functions, Duke Math. J., 30(1963), 51-62.
- [2]. Edwards, J.; A Treatise on Integral Calculus, Chelsea, New York (1954).
- [3]. Erdelyi, A. et. al. ; Higher Transcendental Functions, vol. I, McGraw-Hill, New York (1953).
- [4]. Fox, C.; The G and H function as symmetric Fourier kernels Trans. Amer. Math. Soc. 98 , 1961 , 395-429.
- [5]. Jain, R.N.; Some double integral transformations of certain hypergeometric functions, Math. Japon, 10(1965), 17-26.
- [6]. Rainville, E.D.; Special Functions, Macmillan, New York (1960).
- [7]. Singh, R.P.; A note on double transformation of certain hypergeometric functions, Proc. Edinburgh. Math. Soc. (2),1491965), 221-227.
- [8]. Saxena, R.K.; Some theorems on generalized Laplace transforms, Proc. Nat. Inst. Sc. India, 26 A(1960), 400-413.
- [9]. Srivastava, H.M. and Singhal, J.P.; Double Meijer's transformation of certain hypergeometric functions, Proc. Camb. Phil. Soc. 64(1968).
- [10]. Srivastava, H.M. and Joshi, C.M.; Certain Double Whittaker transforms of generalized hypergeometric functions, The Yokohama Mathematical Journal, vol. XV, (1) (1967), 17-32.
- [11]. Sudland, N., Baumann, B. and Nonnenmacher, T.F.; Open problem: who knows about the Aleph( $\aleph$ )-functions? Frac. Calc. Appl. Annl. 1(4),(1998), 401-402.

IOSR Journal of Engineering (IOSRJEN) is UGC approved Journal with SI. No. 3240, Journal no. 48995.

Yashwant Singh. "On Some Double Integral Transform of Aleph ( $\aleph$ ) -Function." IOSR Journal of Engineering (IOSRJEN), vol. 09, no. 01, 2019, pp. 55-60.