# On contact slant submanifolds in Kenmotsu space forms

## Pradeep Kumar Pandey

Department of Mathematics, Jaypee University of Information Technology, Waknaghat, Solan, Himachal-173234

Abstract: In this paper a sharp estimate of the maximum Ricci curvature function for a special contact slant submanifold M in a Kenmotsu space form  $\overline{M}(c)$ , in terms of the main extrinsic invariant namely the squared mean curvature has been obtained. If submanifold M satisfies the equality case identically, then it is minimal.

Mathematics Subject Classification 2000: 53C40, 53B25, 53D15.

Keywords: Contact slant submanifold, Kenmotsu space form, Mean curvature, Ricci curvature.

#### I. INTRODUCTION

The study of slant submanifold of an almost Hermitian manifold was started by B. Y. Chen ([1], [2]). Further Chen and Tazawa ([3], [4]), furnished examples of slant submanifolds of  $C^2$  and  $C^4$ , while Maeda, Ohnita and Udagawa [5], studied slant submanifolds of Kaehler manifold. The notion of slant submanifold of almost contact metric manifold was introduced by A. Lotta [6]. A class of manifolds was studied and characterized by K. Kenmotsu [7], and the structure so obtained is nowadays known as Kenmotsu structure. In general a Kenmotsu structure is not Sasakian. A Kenmotsu space form is a Kenmotsu manifold with constant  $\phi$ -holomorphic sectional curvature c, denoted by  $\overline{M}(c)$ . Recently, K. Arslan et. al. [8], have established a sharp relationship between the Ricci curvature and the squared mean curvature for submanifolds in Kenmotsu space forms. A (2m+1)-dimensional Riemannian manifold  $\overline{M}$  is said to be an almost contact metric manifold if there exist structure tensors  $(\phi,\xi,\eta,g)$ , where  $\phi$  is a (1, 1) tensor field,  $\xi$  a vector field,  $\eta$  a 1-form and g the Riemannian metric on  $\overline{M}$  satisfying [9]:

- (1.1)  $\phi^2 X = -X + \eta(X)\xi$
- (1.2)  $\phi \xi = 0$
- (1.3)  $\eta(\xi) = 1$
- $(1.4) \qquad \eta(\phi X) = 0$
- $(1.5) g(\phi X, \phi Y) = g(X, Y) \eta(X) \eta(Y)$
- $(1.6) \eta(X) = g(X, \xi)$

for any  $X,Y\in \overline{TM}$ , where  $\overline{TM}$  denotes the Lie algebra of vector fields on  $\overline{M}$ . An almost contact metric manifold  $\overline{M}$  is called a Kenmotsu manifold if [7],

(1.7) 
$$(\overline{\nabla}_X \phi)Y = -g(X, \phi Y)\xi - \eta(Y)\phi X$$
 and  $\overline{\nabla}_X \xi = X - \eta(X)\xi$ 

Where  $\overline{
abla}$  denotes the Levi-Civita connection on  $\overline{M}$  .

The curvature tensor  $\overline{R}$  of a Kenmotsu space form  $\overline{M}(c)$  is given by [7],

(1.8) 
$$\overline{R}(X,Y)Z = \frac{c-3}{4} \{ g(Y,Z)X - g(X,Z)Y \} + \frac{c+1}{4} \{ \eta(X)\eta(Z)Y - \eta(Y)\eta(Z)X + \eta(Y)g(X,Z)\xi - \eta(X)g(Y,Z)\xi - g(\phi X,Z)\phi Y + g(\phi Y,Z)\phi X + 2g(X,\phi Y)\phi Z \}$$

for all  $X, Y, Z \in \overline{TM}$ .

Let M be an n-dimensional submanifold of a Kenmotsu space form  $\overline{M}(c)$  of constant  $\phi$ -sectional curvature c. We denote by  $K(\pi)$  the sectional curvature of M associated with a plane section  $\pi \subseteq T_pM$ ,

 $p \in M$  , and  $\nabla$  the Riemannian connection on M , respectively. We denote by h the second fundamental form of

M and by R and  $\overline{R}$  the curvature tensors of the submanifold M and Kenmotsu space form  $\overline{M}(c)$ , respectively. Then the equation of Gauss is given by

(1.9) 
$$\overline{R}(X,Y,Z,W) = R(X,Y,Z,W) + g(h(X,W),h(Y,Z)) - g(h(X,Z),h(Y,W))$$

for all  $X, Y, Z, W \in TM$ .

Let  $p \in M$  and  $\{e_1, e_2, ..., e_{2m+1}\}$  be an orthonormal basis of the tangent space  $\overline{T_pM}$ , such that  $e_1, e_2, ..., e_n$  are tangent to M at p. Let H denote the mean curvature vector, that is  $H(p) = \frac{1}{n} \sum_{i=1}^n h(e_i, e_i)$ . Now we set

$$(1.10) h_{i,j}^r = g(h(e_i, e_j), e_r), i, j = \{1, 2, ..., n\}, r \in \{(n+1), ..., (2m+1)\}$$

For any vector field X tangent to M, we put  $\phi X = PX + FX$ , where PX and FX denote the tangential and normal components of  $\phi X$  respectively.

## II. CONTACT SLANT SUBMANIFOLDS

K. Arslan et. al. [8] have established a sharp relationship between the Ricci curvature and the squared mean curvature for submanifolds in Kenmotsu space forms. We recall the following theorem for our ready reference.

**Theorem [8]:** Let  $\overline{M}(c)$  be a (2m+1)-dimensional Kenmotsu space form and M an n-dimensional submanifold tangent to  $\xi$ . Then

(i) for each unit vector  $X \in T_n M$  orthogonal to  $\xi$ ,

$$(2.1) Ric(X) \le \frac{1}{4} \left\{ (n-1)(c-3) + \frac{1}{2} \left( 3 \|PX\|^2 - 2 \right) (c+1) + n^2 \|H\|^2 \right\}$$

(ii) A unit tangent vector X orthogonal to  $\xi$  at p satisfies the equality case of (2.1) if and only if

(2.2) 
$$\begin{cases} h(X,Y) = 0, \ \forall Y \in T_p M \ orthogonal \ to \ X \\ 2h(X,X) = nH(p) \end{cases}$$

A submanifold M tangent to  $\xi$  in a Kenmotsu space form  $\overline{M}(c)$  is said to be contact slant submanifold if for any  $p \in M$  and any  $X \in T_pM$  linearly independent on  $\xi$ , the angle between  $\phi X$  and  $T_pM$  is a constant  $\theta$ , called the slant angle of M. A contact slant submanifold which is neither invariant nor anti-invariant is called a proper slant submanifold.

A particular case of contact slant submanifolds, which we call special contact slant submanifolds ([10], [13]). A proper contact  $\theta$ -slant submanifold is said to be special contact slant if

$$(\nabla_{\mathbf{Y}} P) Y = g(Y, PX) \xi - \eta(Y) PX$$

Any 3-dimensional proper contact slant submanifold of a Kenmotsu manifold is a special contact slant submanifold [10].

We denote by  $\mathbf{R}$  the maximum Ricci curvature function on M [11], defined by

$$\mathbf{R}(p) = \max \left\{ Ric(u) \mid u \in T_p^1 M \right\}, \ p \in M$$

where  $T_p^1 M = \{ u \in T_p M \mid g(u, u) = 1 \}.$ 

If n = 3, **R** is the Chen's first invariant  $\delta_M$  introduced in [12]. For n > 3, **R** is the Chen's invariant  $\delta(n-1)$ .

In this section, we establish an inequality for the Chen invariant  $\, R \,$  and show that any special contact slant submanifold which satisfies the equality case identically is minimal. Now we show

**Theorem (2.1):** Let M be an (n+1)-dimensional special contact slant submanifold in a (2n+1)-dimensional Kenmotsu space form  $\overline{M}(c)$  of constant  $\phi$ -sectional curvature c. Then for each unit vector X orthogonal to  $\mathcal{E}$ , we have

(2.3) 
$$\mathbf{R} \leq \frac{1}{4} \{ (n+1)^2 \|\mathbf{H}\|^2 + \mathbf{n}(c-3) + \frac{1}{2} (3\cos^2\theta - 2)(c+1) \}$$

If M satisfies the equality case in (2.3) identically, then M is a minimal submanifold.

**Proof:** The inequality (2.3) is an immediate consequence of the inequality (2.1). To ascertain minimality, we assume that M is a special contact slant submanifold of Kenmotsu space form  $\overline{M}(c)$ , which satisfies the equality case of (2.3) at a point  $p \in M$ . Now without loss of generality we may choose an orthonormal basis  $\{e_0 = \xi, e_1, \ldots, e_n\}$  of  $T_pM$  such that  $\mathbf{R}(p) = Ric(e_n)$ . Then by (2.2) it follows that

(2.4) 
$$h(e_i, e_n) = 0, \forall i \in \{0, ..., n-1\}; h(e_n, e_n) = \sum_{i=1}^{n-1} h(e_i, e_i)$$

where  $\{e_0=\xi\,,e_1\,,...,e_n\,,e_{n+1}\,,...,e_{2n}\}$  is the orthonormal basis with  $e_{n+j}=(\sec\theta)\,Pe_j$  ,  $j\in\{1,2,...,n\}$  and  $e_{n+1}$  parallel to the mean curvature vector H(p) .

Let A denote the Weingarten map of M in  $\overline{M}(c)$ . Then it is easy to show that  $A_{\phi X}Y = A_{\phi Y}X$  for all vector fields X,Y tangent to M (see [10]). Hence we have  $h_{i,j}^{n+k} = h_{i,j}^{n+j} = h_{i,j}^{n+i}$  for any  $i,j,k \in \{1,2,\ldots,n\}$ . Now using equations (2.4) a simple computation yields H(p) = 0, therefore M is a minimal submanifold.

We have the following characterization of n-dimensional special slant submanifolds in the sphere  $S^{2n+1}$  which satisfy the equality case of (2.3) identically.

**Theorem (2.2):** Let M be an (n+1)-dimensional special contact slant submanifold of the (2n+1)-dimensional sphere  $S^{2n+1}$ . Then M satisfies the equality case of (2.3) identically if and only if M is minimal and ruled. **Proof:** The theorem (2.1) implies that M is a minimal submanifold. From the theorem [8], it is obvious that the equality case of (2.3) holds identically if and only if the relative null space

$$\mathbf{N}_{p} = \{X \in T_{p}M \mid h(X,Y) = 0, \forall Y \in T_{p}M\} \neq \{0\}$$

at any  $p \in M$ .

Let  $U_l$  denotes interior of the subset consisting of points in M such that the relative null space at p has dimension l. It follows that for some l,  $U_l \neq \varphi$ , where  $l \in \{1,...,n\}$ .

Let  $X, Y \in \mathcal{N}$ , and Z any vector field tangent to M. By applying Codazzi equation

$$(\overline{\nabla}_{_{\!Y}}h)(Y,Z) = (\overline{\nabla}_{_{\!\!Z}}h)(X,Y),$$

we get  $h(\nabla_X Y, Z) = 0$ . Then N is integrable on  $U_l$  and each leaf of  $N \mid U_l$  is an l-dimensional totally geodesic submanifold of  $S^{2n+1}$ . Thus submanifold M contains a geodesic of  $S^{2n+1}$  through each point  $p \in U_l$ .

Since M is the union of the closure of all  $U_l$ , we conclude by continuity that M contains a geodesic of the ambient space through each point in M. Hence, M is minimal and ruled. The converse follows from straightforward computation.

### REFERENCES

- 1). B. Y. Chen, Slant immersions, Bulletin of the Australian Mathematical Society, 41, 1990, 135-147.
- 2). B. Y. Chen, Geometry of slant submanifolds (Katholieke Universiteit. Leuven, 1990).
- 3). B. Y. Chen, and Y. Tazawa, Slant submanifolds in complex Euclidean spaces, *Tokyo Journal of Mathematics*, 14 (1), 1991, 101-120.
- 4). B. Y. Chen, and Y. Tazawa, Slant surfaces with codimension 2, *Ann. Fac. Sci. Toulouse Math. XI* (3), 1990, 29-43.
- 5). S. Maeda, Y. Ohnita, and S. Udagawa, On slant immersions into Kaehler manifolds, *Kodai Mathematical Journal*, 16, 1993, 205-219.
- 6). A. Lotta, Slant submanifolds in contact geometry, Bull. Math. Soc. Roumanie, 39, 1996, 183-198.
- 7). K. Kenmotsu, A class of almost contact Riemannian manifolds, *Tohoku Mathematical Journal*, 21, 1972, 93-103.
- 8). K. Arslan, R. Ezentas, I. Mihai, C. Murathan and C. Ozgür, Ricci curvature of submanifolds in Kenmotsu space forms, *IJMMS* 29 (12), 2002, 719-726.
- 9). D. E. Blair, *Contact manifolds in Riemannian geometry* (Lecture notes in Mathematics, Springer Verlag 509, 1976, Berlin-New York).
- 10). R. S. Gupta, S. M. Khursheed and M. H. Shahid, Slant submanifolds of a Kenmotsu manifold, *Radovi Matematicki* 12, 2004, 205-214.
- 11). B. Y. Chen, On Ricci curvature of isotropic and Lagrangian submanifolds in complex space forms, *Archiv. Math.*, 74, 2000, 154-160.
- 12). B. Y. Chen, some pinching and classification theorems for minimal submanifolds, *Arch. Math.*, 60, 1993, 568-578.
- 13). I. Mihai, and V. Ghisoiu, Minimality of certain contact slant submanifolds in Sasakian space forms, *Int. J. Pure and Appl. Math. Sci.*, 1, 2004, 95-99.